
MYCOTAXON

Volume 116, pp. 121–124

April–June 2011

DOI: 10.5248/116.121

***Ileodictyon gracile*, new to Italy**

ALESSANDRO SAITTA, RICCARDO COMPAGNO, ALFONSO LA ROSA,
DANILO LOPEZ & GIUSEPPE VENTURELLA*

*Dipartimento di Biologia vegetale e Biodiversità, Università di Palermo,
Via Archirafi 38, I-90123 Palermo, Italy*

* CORRESPONDENCE TO: gvent@unipa.it

ABSTRACT — The first record of *Ileodictyon gracile* from Italy is reported, accompanied by notes on the taxonomy, ecology, and distribution of this rare gasteromycete.

KEY WORDS — *Phallaceae*, rare Mediterranean fungi, gasteromycetoid fungi

Introduction

Gasteromycetoid fungi encompass terrestrial, epigeous and hypogeous and saprobic and ectomycorrhizal species. They are cosmopolitan but are more frequently found in warm, dry habitats (Kirk et al. 2008). Modern accounts of this interesting but taxonomically complex group is particularly limited in Italy where after the monograph by Petri (1909), only Sarasini (2005) has written on epigeous gasteromycetes in the country. In the absence of funding for studies of gasteromycetoid fungi, reports of undescribed species or new records from Italy are usually restricted to an occasional observation and/or generalized papers devoted to fungal diversity. This is the first record of *Ileodictyon gracile* in Italy, recently collected by us on the island of Pantelleria, Egadi Archipelago in Trapani province in south-eastern Sicily.

Materials & methods

Basidiomata were identified while fresh and microscopic features were observed in H₂O using a Leica microscope DMLB; spore measurements were based on 50 observations. Nomenclature follows Index Fungorum (<http://www.indexfungorum.org/Names/Names.asp>). The description is based both on personal observation, augmented by characters cited in Sarasini (2005) and Bougher & Syme (1998). The collection is curated in the fungal dried reference collection of the Herbarium Mediterraneum Panormitanum, Palermo (PAL).

Taxonomy

Ileodictyon gracile Berk., London J. Bot. 4: 69. 1845.

FIG. 1

Unripe basidiomata, sessile, irregularly rupturing at apex, egg-shaped, globose, subglobose or sphaerical, consistency jelly-like, 1.5 cm in diam. Peridium white, with long white unbranched rhizomorphs emanating from various points. Ripe basidiomata, sessile, with receptacle (lattice) up to 80–200 mm broad, white or cream, sphaerical or broadly ellipsoidal, sessile in a ruptured, white, membranous volva. Lattice of isodiametric-polygonal interspaces and flattened, firm struts that are grooved but not broadened at junctions and have glebal slime adhering to inside surface. Peridium in two layers with hyaline hyphae, 3.5–6.5 wide: exoperidium, white, mat, thin and membranous; endoperidium thick and gelatinous. Gleba olive-brown. Basal mycelium of white mycelia strands. Odour foetid or like sour milk. Spores hyaline, (4–)4.5–5.5(–6) × 1.8–2.4 µm, narrowly ellipsoidal, smooth, thin-walled. Basidia clavate, 15–25 × 4–6 µm. Cystidia absent.

SPECIMEN EXAMINED: ITALY. SICILY: province of Trapani, Egadi archipelago, Island of Pantelleria, Bugeber, 200 m, wood of *Quercus ilex* L. (*Fagaceae*) with shrubs of *Cistus salviifolius* L. (*Cistaceae*) in the ground cover, 2 Jan 2009, coll. R. Compagno, A. La Rosa & D. Lopez (PAL 00001/2011).

COMMENTS — Hosaka et al. (2006) recently reclassified gomphoid-phalloid fungi based on molecular phylogenetic characters, and Kirk et al. (2008) included the genus *Ileodictyon* Tul. & C. Tul. (*Phallaceae* Corda) in the *Phallales* E. Fisch. Berkeley (1845) separated the genus *Ileodictyon* from *Clathrus* P. Micheli ex L. principally based on possession of tubular (not cellular) ribs and a different volval morphology. Dring & Rose (1977) found that the virtually sphaeroidal symmetry further differentiated *I. gracile* and *I. cibarium* Tul. & C. Tul. from *Clathrus* s. str. *Ileodictyon gracile* is separated from *I. cibarium* based on morphological differences of the receptacle, intersections, and creases (Dring 1980).

Discussion

The so-called lattice or basket fungus is a saprotrophic gasteromycete usually collected on ground, commonly on the edge of tracks in forests and urban areas (Bougher & Syme 1998). *Ileodictyon gracile* has been frequently reported from Australia (Berkeley & Broome 1879, Cunningham 1944, Bougher & Syme 1998); May et al. 2003) also provide color and black and white illustrations of basidiomata and microscopic characters. The species also occurs in New Zeland, Samoa, Japan, Korea, and Africa (Calonge & Romero Zarco 1989, Dring & Rose 1977, Ka et al. 2004, Malençon & Bertault 1970, Pennycook & Galloway 2004). European distribution is restricted to Spain (Calonge & Romero Zarco




FIG. 1. Basidiomata of *Ileodictyon gracile*.

1989), Canary Islands (Chávez Barreto & Escobio García 2009), and Portugal (Pinto-Lópes 1942, 1944; Louro et al. 2009).

We found that *Ileodictyon gracile* shares similar habitats and phenology (winter) in Portugal, Spain, and Sicily. In Portugal it was collected under *Cistus ladanifer* L. in a *Quercus rotundifolia* Lam. forest, in Spain it was recorded in *Pinus pinea* L. woods with *Cistus salviifolius* as ground cover over sandy soil (Calonge & Romero Zarco 1989), and in Sicily in a *Q. ilex* wood with *C. salviifolius* shrubs. Other reported habitats include *Eucalyptus* reforestations (Chávez Barreto & Escobio García 2009), open ground near forest trails and urban areas (Bougher & Syme 1998), and on sandy soil in glades, woods, parks, and gardens (Sarasini 2005).

Our report represents the southernmost limit of the European distribution of *I. gracile*.

Acknowledgements

The authors wish to thank Dr Georgios I. Zervakis (Greece) and Dr Alfredo Vizzini (Italy) for critically reviewing the manuscript.

Literature cited

- Berkeley MJ. 1845. Decades of fungi. Dec. III.-VII. Australian fungi. London Journal of Botany 4: 42-73.
- Berkeley MJ, Broome CE. 1879. List of fungi from Brisbane, Queensland; with description of new species. Transactions of the Linnean Society of London, Botany, ser. 2, 1(6): 399-408.
- Bougher NL, Syme K. 1998. Fungi of southern Australia. Nedlands, University of Western Australia Press. 391 p.
- Calonge FD, Romero Zarco C. 1989. *Ileodictyon gracile* Berk. (*Gasteromycetes*), nuevo para el catálogo micológico español. Boletín Sociedad Micológica de Madrid 13: 171-174.
- Chávez Barreto D, Escobio García VJ. 2009. Algunas nuevas citas para el Catálogo fúngico de las islas de Gran Canaria y Tenerife, (Islas Canarias). Cantarela, Boletín de la Sociedad Micológica de Gran Canaria 44: 1-4.
- Cunningham GH. 1944. The gasteromycetes of Australia and New Zealand. Dunedin, John McIndoe. 236 p.
- Dring DM. 1980. Contributions towards a rational arrangement of the *Clathraceae*. Kew Bulletin 35(1): 1-96. doi: [10.2307/4117008](https://doi.org/10.2307/4117008)
- Dring DM, Rose AC. 1977. Additions to West African phalloid fungi. Kew Bulletin 31(3): 741-751. doi: [10.2307/4119427](https://doi.org/10.2307/4119427)
- Hosaka K, Bates ST, Beever RE, Castellano MA, Colgan III W, Domínguez LS, Nouhra ER, Geml J, Giachini AJ, Kenney SR, Simpson NB, Spatafora JW, Trappe JM. 2006. Molecular phylogenetics of the gomphoid-phalloid fungi with an establishment of the new subclass *Phallomycetidae* and two new orders. Mycologia 98(6): 949-959. doi: [10.3852/mycologia.98.6.949](https://doi.org/10.3852/mycologia.98.6.949)
- Ka K-H, Hur T-C, Ryoo, C-I. 2004. A note on *Ileodictyon* (*Clathraceae*) in Korea. Mycobiology 32(1): 54-56. doi: [10.4489/MYCO.2004.32.1.054](https://doi.org/10.4489/MYCO.2004.32.1.054)
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Dictionary of the fungi. 10th Edition. Wallingford, CAB International. 771 p.
- Louro R, Calado M, Pinto B, Santos-Silva C. 2009. Epigeous macrofungi of the Parque de Natureza de Noudar in Alentejo (Portugal). Mycotaxon 107: 49-52.
- Malençon G, Bertault R. 1970. Flore de champignons supérieurs du Maroc. Tome I. Rabat, Faculté de Science. 601 p.
- May TW, Milne J, Shingles S, Jones RH. 2003. Fungi of Australia, Vol. 2B. Catalogue and bibliography of Australian fungi 2. *Basidiomycota* p.p. & *Myxomycota*. Canberra, ABRS/CSIRO Australia. 452 p.
- Pennycook SR, Galloway D.J. 2004. Checklist of New Zealand "fungi". 401-488, in: EHC McKenzie (ed.). Introduction to fungi of New Zealand. Fungi of New Zealand: Ngā Harore o Aotearoa, Vol. 1. Hong Kong, Fungal Diversity Press.
- Petri L. 1909. *Fungi. Gasterales. Secotiaceae, Lycoperdaceae, Sclerodermataceae*. Flora Italica Cryptogama, 1(5). Rocca S. Casciano, Stabilimento Tipografico Cappelli. 140 p.
- Pinto-López J. 1942. Contribuição para o estudo dos Basidiomicetas portugueses, II. Boletim da Sociedade Broteriana, Ser. 2, 16: 215-225.
- Pinto-López J. 1944. Contribuição para o estudo dos Basidiomicetas portugueses. Brotéria, Ciências Naturais 13: 230-238.
- Sarasini M. 2005. Gasteromiceti epigei. Trento/Vicenza, Associazione Micologica Bresadola/Fondazione Centro Studi Micologici dell'AMB. 406 p.